



## Prevalence of Geohelminth Parasites of Horses in Dutsinma Metropolis, Katsina State Nigeria

J. B. Orpin<sup>1\*</sup>, I. Mzungu<sup>1</sup> and M. S. Hussaini<sup>1</sup>

<sup>1</sup>Department of Biological Sciences, Federal University, Dutsinma, PMB 5001, Dutsinma, Katsina State, Nigeria.

### Authors' contributions

This work was carried out in collaboration between all authors. Author JBO designed the study and performed the statistical analysis of the manuscript. Author IM managed the analyses and the discussion of the study and Author MSH managed the literature searches and the introduction. All authors read and approved the final manuscript.

### Article Information

DOI: 10.9734/JABB/2018/v20i230072

#### Editor(s):

- (1) Dr. Jamal M. Abo Omar, Professor, Department of Animal Production, An-Najah National University, Palestine.
- (2) Dr. George Perry, Dean and Professor of Biology, University of Texas at San Antonio, USA.
- (3) Cosmas Nathanailides, Dept. Fisheries and Aquaculture Technology, Technological Educational Institute of West Greece, Greece.

#### Reviewers:

- (1) Hasibe Cingilli Vural, Necmettin Erbakan University, Turkey.
  - (2) Roberto Amerigo Papini, Università di Pisa, Italy.
  - (3) Ukwubile, Cletus Anes, Federal Polytechnic Bali, Nigeria.
- Complete Peer review History: <http://www.sdiarticle3.com/review-history/34737>

Original Research Article

Received 28 June 2017  
Accepted 03 September 2017  
Published 01 March 2019

### ABSTRACT

This study was carried out to investigate the prevalence of geohelminth parasites of horses in Dutsinma metropolis between May to August, 2016. Fresh fecal sample were obtained from the rectum of 48 randomly selected horses. A floatation technique was used for the lab work experiment. From the result, out of the total sample collected 6 (12.50%) were positive with eggs of two parasites Strongyles and Oxyuris spp with the prevalence of (6.25%) for each. Among the three points of sample collection, the highest prevalence was recorded at Kadangaru (20.00%) followed by Unguwa Yandaka (18.18%) and no case recorded at Hayingada.  $X^2$  cal = 36.84 at  $P=0.05$  and 2df. The highest prevalence was recorded among the males (18.52%) while only 1 female was affected with the prevalence of (4.76%)  $X^2$  cal = 36.57 at  $P=0.05$  and 2df. Prevalence was recorded in adult of 14.29% while young horses recorded a prevalence of 10.00%,  $X^2$  cal = 36.84 at  $P=0.05$  Strongyles and Oxyuris spp were recognized as important geohelminth parasites in Horses in Dutsinma Local Government Area of Katsina State with various rates of infection (12.50%). Managerial control tools should be integrated to improve prevention of geohelminth parasites by communication, information and health education.

\*Corresponding author: E-mail: jorpin@fudutsinma.edu.ng;

**Keywords:** Prevalence; faecal samples; geohelminth; parasites; horses.

## 1. INTRODUCTION

Geohelminth parasites are soil-transmitted helminths which are groups of parasitic worms causing human and other Animals infection through contact with parasite eggs or larvae that thrive in the warm and moist soil of the world's tropical and subtropical countries. As adult worms, the soil-transmitted helminths live for years in the human gastrointestinal tract. More than a billion people are infected with at least one species of geohelminth parasites worldwide. They are considered together because it is common for a single individual, especially a child living in a less developed country, to be chronically infected with all three worms. Such children have malnutrition, growth stunting, intellectual retardation, and cognitive and educational deficits.

Parasitic diseases are the major obstacle in the growth and development of animal health all over the world. Horses, among most domestic animals are reported to be more susceptible to large number of parasites and may harbour different species at any time. An apparently healthy horse can harbour over one half million geohelminth parasites, trematodes, cestodes and nematodes [1]. This is because, the gastrointestinal tract provides favorable environment for the survival and proliferation of many of these parasites.

Horses (*Equus ferus caballus*) are one of the two extant sub species of horse *Equus ferus* [2]. They evolve over 45-55 million years. Humans began to domesticate horse around 4000 B C. Horses and humans interact in a wide variety of sport competition and man competitive activities.

Intestinal parasites such as helminths usually produce insidious diseases in animals. Infected horse may show signs of weakness, emaciation, restlessness, unthriftiness, diarrhea, anemia and sometimes intestinal obstruction or perforation.

The aim of this study is to determine the Prevalence of geohelminth Parasites of Horses in Dutsinma Metropolis, Katsina State with the following objectives:

- To determine the prevalence of geohelminth parasites of horses in Dutsinma metropolis.
- To identify the geohelminth parasites among horses in the study area.

- To isolate the geohelminth parasites species in the Horses.

## 2. STUDIED AREA

Dutsinma (Lat 12°27'01.18" N/ Long 7°29'.29" E), Katsina State, Nigeria Dutsinma Local Government covers an area of 527 km squares and has a population of 16971, at the 2006 census. Total annual rainfall around Dutsinma is about 800 mm Retrieved 10<sup>th</sup> July 2011 from www.albarkablog.com. The inhabitants of the local government are predominantly Hausa and Fulani by tribe. Their main occupations are farming and animal rearing.

### 2.1 Sample Population

A total of 48 faecal samples of horses were collected from different places within Dutsinma metropolis including Unguwar Yandaka, Hayin gada, and Kadangaru areas.

### 2.2 Sample Collection

Samples were collected between the periods of May to August, 2016. Faecal samples were randomly collected from the rectum of horses in each place of the study area during the early hours of the day while owners restrained the animals [3]. Aseptic measures were maintained during collection. Each sample was collected into a clean polythene bag while wearing gloves, labeled, noting the sex and age of the animal. These samples were immediately conveyed to the Biology Laboratory of Department of Biological Science, Federal University Dutsinma for examination.

### 2.3 Sample Examination

All faecal samples collected from the rectum of horses were screened eggs/oocysts of the parasite using the floatation technique. A technique with a principle where the floatation medium (Saturated salt NaCl solution) aids the rise of eggs and/or oocysts of geohelminth parasites by virtue of the medium's high specific gravity, allowing the faecal material to settle to the bottom.

### 2.4 Preparation of Floatation Medium

To prepare the floatation medium, 10 grams of NaCl was dissolved in 500 mL distilled water







Fig. 2. Map of the Study Area showing the



**Fig. 2. Map of Dutsinma**  
 Source:- Cnes/Spot Image 2014

**Table 1. Prevalence of geohelminth parasites in the study area**

Location	No. screened	No. positive	Prevalence	E	X <sup>2</sup>
Hayin Gada	16	0	0.00	0.00	0.00
Unguwar Yandaka	22	4	18.18	0.50	24.50
Kadangaru	10	2	20.00	0.25	12.25
Total	48	6	12.50	0.75	36.75

X<sup>2</sup> cal = 36.75 > X<sup>2</sup> tab = 10.60, therefore significant difference exist at P ≤ 0.05 and 2df.  
 KEY: X<sup>2</sup> = Chi-square, and E = expected value

**3.2 Prevalence of Geohelminth Parasites in Relation to Sex**

Out of the 48 horses examined, 27 were males and 21 were females. The highest prevalence was recorded among the males (18.52%) while only 1 female was affected with the prevalence of (4.76%).

**3.3 Prevalence of Geohelminth Parasites in Relation to Age**

28 (58.33%) adult and 20 (41.67%) young Horses were examined. Slightly highest prevalence was recorded at adult with the prevalence of 14.29% while low prevalence was founded at young horses with the prevalence of 10.00%.

**Table 2. Prevalence of geohelminth parasites in relation to the sex of horses**

Location	Male					Female				
	No. scre.	Positive	%	E	X <sup>2</sup>	No. scre.	Positive	%	E	X <sup>2</sup>
Unguwar Yandaka	12	4	33.33	0.74	14.17	10	0	0.00	0.00	0.00
Hayin gada	9	0	0.00	0.00	3.45	7	0	0.00	0.00	18.05
Kadangaru	6	1	16.67	0.19	0.00	4	1	25.00	0.05	0.00
Total	27	5	18.52		17.62	21	1	4.76	0.05	18.05

$X^2_{cal} = 36.57 > X^2_{tab} = 10.60$ , therefore significant difference exist at  $P \leq 0.05$  and 2df.

Key: X<sup>2</sup>=Chi-square, E = expected value, and % = prevalence and No = number screened

**Table 3. Prevalence of geohelminth parasites in relation to the age of horses**

Location	Adult					Young				
	Adult	Positive	%	E	X <sup>2</sup>	Young	Positive	%	E	X <sup>2</sup>
Unguwar Yandaka	13	3	23.08	0.43	15.36	9	1	11.11	0.10	8.10
Hayin gada	9	0	0.00	0.00	0.00	7	0	0.00	0.00	0.00
Kadangaru	6	1	16.67	0.14	5.28	4	1	25.00	0.10	8.10
Total	28	4	14.29	0.57	20.64	20	2	10.00	0.20	16.20

$X^2_{cal} = 36.84 > X^2_{tab} = 10.60$ , therefore significant difference exist at  $P \leq 0.05$  and 2df

Key: X<sup>2</sup>=Chi-square, E = expected value, and % = prevalence

**Table 4. Quantitative prevalence of geohelminth parasite identified in the study area**

Location	Strongyles					Oxyuris spp				
	No. screened	Positive	%	E	X <sup>2</sup>	No. screened	Positive	%	E	X <sup>2</sup>
Unguwar Yandaka	22	1	4.55	0.06	14.73	22	2	9.09	0.13	<b>26.90</b>
Hayin gada	16	0	0.00	0.00	0.00	16	0	0.00	0.00	0.00
Kadangaru	10	2	20.0	0.13	26.90	10	1	10.00	0.06	14.73
Total	48	3	6.25	0.19	41.63	48	3	6.25	0.19	41.63

$X^2_{cal} = 83.26$  is significant difference exist at  $P \leq 0.05$  and 2df.

Key: X<sup>2</sup>=Chi-square, E = expected value, and % = prevalence

### 3.4 Geohelminth Parasites Identified in the Study Area

Eggs of the parasites as identified in the study area including those of *Strongyles* and *Oxyuris spp* recorded the same prevalence of (6.25%) each.

## 4. DISCUSSION

This study was designed to investigate the Prevalence of Geohelminth Parasites of Horses in Dutsinma. A total of 48 Horses from Unguwar Yandaka, Hayingada and Kadangaru were screened for the research, eggs of two species of geohelminth parasites were recorded 12.50% (*Strongyles* and *Oxyuris spp*). This agrees with previous studies by [6,7] which further explained the predomination of Strongyle eggs with prevalence rates of 58.5%, 57.14%, and 66.67%, respectively. The overall geohelminth prevalence appears to be low, in which only 6(12.50%) were harboring one or more Geohelminth parasites.

Among the three point of sample collection, the highest prevalence was recorded at Kadangaru (20.00%) followed Unguwa Yandaka (18.18%) and no case recorded at Hayingada, Table 1. This is low when compared to that of studies in Ethiopia [8] estimating prevalence in horses to be 91%, and similar estimates of 80% [9] and 81% [10] in Mexico. Potential variation arising from management differences and differences with respect to topography, climate, animal working conditions and seasonality make direct comparisons difficult. The differences in intensity of infection could be due to variations in parasite biology relating to climatic conditions, pasture infection intensity relating to grazing practices and/or differences in use of anthelmintics; detailed information on such variables would be required to determine their importance.

Out of the 48 horses examined, 27 were males and 21 were female horses. The highest prevalence was recorded among the males (18.52%) while only 1 female was affected with

the prevalence of (4.76%). This difference in the prevalence is statistically significant ( $P < 0.05$ ), Table 2. Which is in line with the work of [11] who reported a higher prevalence of (22.09%) male than that of the female Horses (13.56%).

28 (58.33%) adult and 20 (41.67%) young Horses were examined. Slightly highest prevalence was recorded at adult with the prevalence of 14.29% while low prevalence was founded at young horses with the prevalence of 10.00%, Table 3. This is not in line the work of [12] who recorded a slightly higher rate among young (100%) horses than adult (97.8%). This might be associated with apparent inability of the adult younger age groups to develop adequate acquired immunity predisposing them to high risks of severe infection with geohelminth parasites when compared with adult donkeys. Higher infection rates and more severe infections reflect lack of immunity in younger population [13].

The low prevalence we found (12.50% of horses) is similar to the findings of the 2009 survey from Greece, that found *Strongyle* eggs in 25.6% of horses [14]. But this finding is less than the 78% prevalence that was found in the 1986 survey [15]. It might be due to routine deworming programs that might were introduced to most of the horse population in this study area. This decline in prevalence was statistically significant ( $P < 0.05$ ), it may suggest a trend. This concurs with another study from Louisiana conducted over similar time scale, which demonstrated a decrease in the infestation of most geohelminth species [16].

However, the reasons why the parasites vary between age and sex groups is require further investigation, base on my knowledge this is the first attempt to investigate the geohelminth parasites in Dutsinma metropolis. Such further studies could encourage more specific needs for owner education that could reduce endo-parasites burdens and also improve horse welfare.

*Oxyuris equi* with prevalence rate of 6.25 % was very low when compared with the work of [15,8].

## 5. CONCLUSION

It has been deduced in this research that *Strongyles* and *Oxyuris spp* were identified as important geohelminth parasites in Horses in Dutsinma Local Government Area of Katsina State.

## 6. RECOMENDATION

- ❖ Based on the results obtained and in order to avoid frequent infection, it is recommended that routine feacal samples go through Parasitological diagnostic techniques to determine cases of geohelminth parasites.
- ❖ Communication information and health education be integrated as preventive tool for horse owners.
- ❖ Owners of horses should routinely take them to Veterinary Clinics for accurate and appropriate diagnosis.

## ACKNOWLEDGEMENT

The authors would like to acknowledge all staff of Federal University Dutsinma and all the reviewers of this paper.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Martins S, Krecek RC, Milne SA. Prevalence and biodiversity of helminth parasites in donkeys from South Africa. *Journal of Parasitology*. 2009;86:756-762.
2. Getachew AM, Trawford AF, Feseha G, Reid SWJ. Gastrointestinal parasites of working donkeys in Ethiopia. *Tropical Animal Health and Production*. 2010;42: 27-33.
3. Abbott EM. Larval cyathostomosis, In: Part 1 *Equine Practice*. 2008;20(3):56 – 67.
4. Jorgen S, Brain R. Isolation of infection. *Dictyocaulus* larvae from herbage. *Vet. Parasitology*. 1994;16(18):16.
5. Yagoob G. Seroepidemiology of *Neospora* Sp. In horses in East Azerbaijan Province of Iran, *Research Journal of Biological Sciences*. 2015;6 (5);224-226.
6. Saeed K, Qadir Z, Ashraf K, Ahmad N. Role of intrinsic and extrinsic epidemiological factors on strongylosis in horses. *Journal of Animal and Plant Sciences*. 2010;20:277-280  
ISSN: 1018-7081.
7. Wannas HY, Dawood KA, Gassem GA. Prevalence of gastro-intestinal parasites in

- horses condition of horses and donkeys in Al Diwaniyah Governorate, AL-Qadisiya Journal of Vet. Med. Sci. 2012;11(1):2012.
8. Fikru R, Bizunesh T. Prevalence of equine gastrointestinal parasites in western highlands of Oromia, Ethiopia, Bulletin of Animal Health and Production in Africa. 2005;53:161-166.
  9. Burden FA, du Toit N, Hernandez-Gil N, Prado-Ortiz O, Trawford AF. Selected health and management issues for working donkeys in rural Mexico, Tropical Animal Health and Production. 2010;42:597-605.
  10. du Toit N, Crewe W. Haemoparasites. In: Blacklock and Southwell, A guide to human parasitology, Laboratory Techniques, 10<sup>th</sup> Edition, H.K. Lewis and Co Ltd, ELBS series. 2008;177-186.
  11. Hinney B, Wirtherle NC, Kyule M, Miethe N, Zessin KH, Klausen PH. Prevalence of helminths in horses in the state of Brandenburg, Germany, Parasitol Res. 2011;108:1083-1091.
  12. Morariu S, Bogdan AT, Dărăbuș G. Heminth parasites in horses from ten locations of Timiș County, Bulletin of University of Agricultural Sciences and Veterinary Medicine Cluj-Napoca. 2012; 69(1/2):381-384.
  13. Urquhart GM, Aremour JV, Duncan AM, Dunn FW, Jennis FW. Veterinary parasitology 2 edition. The University of Glasgow, Black Well Science, Scotland. 1996;3-137.
  14. Papazahariadou M, Papadopoulos E, Diakou A, Ptochos S. Gastrointestinal parasites of stabled and grazing horses in Central and Northern Greece. Journal of Equine Veterinary Science. 2009;29(4): 233-236.
  15. Crane M, Khallaayoune K, Christley R, Scantlebury C, Faoud M. The effect of an anthelmintic programme for working equids in Morocco: A comparison of treated and placebo control Populations. In: Proceedings of the 9<sup>th</sup> International Equine Colic Research, Liverpool, UK. 2008;171-172.
  16. Ayele CF. Principal health problems of donkeys in Dugda Bora district of Ethiopia. In: Proceedings of the 5<sup>th</sup> International Colloquium on Working Equines. Sidmouth. 2006;162-168.

© 2018 Orpin et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<http://www.sdiarticle3.com/review-history/34737>